arXiv:1112.1488v1 [math.PR] 7 Dec 2011

The Skitovich-Darmois theorem for discrete and compact totally disconnected Abelian groups

I.P. Mazur

December 8, 2011

1 Introduction

The classical Skitovich-Darmois theorem states ([14],[1], see also [11, Ch. 3]): Let $\xi_i, i = 1, 2, \ldots, n$, $n \geq 2$, be independent random variables, and α_i, β_i be nonzero numbers. Suppose that the linear forms $L_1 = \alpha_1\xi_1 + \cdots + \alpha_n\xi_n$ $L_2 = \beta_1\xi_1 + \cdots + \beta_n\xi_n$ are independent. Then all random variables ξ_i are Gaussian.

The Skitovich-Darmois theorem was generalized into various algebraic structures, in particular, into locally compact Abelian groups([3]-[9],[10],[12]). In these researches the random variables take values in a locally compact Abelian group X, coefficients of the linear forms are topological automorphisms of X, and the number of the linear forms are two. In the article we continue these researches and study the Skitovich-Darmois theorem in the case when random variables take values in different classes of locally compact Abelian groups but the number of linear forms more than 2.

Throughout the article X is a second countable locally compact Abelian group. Let $Aut(X)$ be the group of topological automorphisms of X, $\mathbb{Z}(k) = \{0, 1, 2 \ldots, k-1\}$ be the group of residue modulo k. Let $x \in X$. Denote by E_x the degenerate distribution, concentrated at the point x. If K is a compact subgroup of X, denote by m_K the Haar distribution on K. Denote by $I(X)$ the set of shifts of such distributions, i.e. the distributions of the form $m_K * E_x$, where K is a compact subgroup of X, $x \in X$.

The first step to generalize the Skitovich-Darmois theorem on locally compact Abelian groups is to consider the case of finite Abelian groups. This problem was studied first in [5]. Note that the idempotent distributions on a finite Abelian group can be regarded as analogues of the Gaussian distributions on real line. Let $\xi_i, i = 1, 2, \dots, n, n \geq 2$, be independent random variables with values in a finite Abelian group X and distributions μ_i. Let α_j, β_j be automorphisms of X. Consider the linear forms $L_1 = \alpha_1\xi_1 + \cdots + \alpha_n\xi_n \quad L_2 = \beta_1\xi_1 + \cdots + \beta_n\xi_n$. G.M. Feldman proved that the class of groups, on which the independence of $L_1 \quad L_2$ implies that all $\mu_i \in I(X)$, are poor and consists of the groups of the form ([5])

$$\mathbb{Z}(2^{m_1}) \times \cdots \times \mathbb{Z}(2^{m_l}), 0 \leq m_1 < \cdots < m_l. \tag{1}$$

On the other hand if we consider n linear forms of n independent random variables, then the Skitovich-Darmois theorem is valid for an arbitrary finite Ableian group. Namely, the following theorem holds ([12]):

Theorem 1.1 *Let $\xi_i, i = 1, 2, \dots, n, n \geq 2$, be independent random variables with values in a finite Abelian group X and distributions μ_i. If the linear forms $L_j = \sum_{i=1}^n \alpha_{ij}\xi_i$, where $\alpha_{ij} \in Aut(X), i, j = 1, 2, \dots, n$, are independent, then $\mu_i \in I(X), i = 1, 2, \dots, n$.*

Our first aim is to generalize Theorem 1.1 into discrete groups. Using this result we extend Theorem 1.1 into groups of the form $\mathbb{R}^m \times D \times K$, where D is a discrete, K is a compact totally disconnected group of the special form, $m \geq 0$. Finally we describe conditions on compact totally disconnected group, which are necessary and sufficient for the validity of the Theorem 1.1.

2 Definitions and designations

Let X be a second countable locally compact Abelian group. Let $\{X_\lambda : \lambda \in \Lambda\}$ be a non empty family of groups. Denote by $\mathbf{P}_{\lambda\in\Lambda}X_\lambda$ the direct product of the groups X_λ. If $X = X_\lambda$ for all $\lambda \in \Lambda$, then the direct product of the groups X_λ we denote by $X^{\mathfrak{n}}$, where $\mathfrak{n}$ is a cardinal number of Λ.

Let p be a prime number. Denote by $\mathfrak{P}$ the set of all prime number. Denote by $\triangle_p$ the group of p-adic integers. We recall that an Abelian group is called a p-primary if every nonzero element of this group has order p^m for some natural m.

Let μ be a distribution on X. The characteristic function of the distribution μ is defined by the formula ([2])

$$\hat{\mu}(y) = \int_X (x, y) d\mu(x), y \in Y.$$

Recall ([13]), that a distribution μ on X is called Gaussian, if its characteristic function can be represented in the form

$$\hat{\mu}(y) = (x, y) \exp\{-\phi(y)\}, \quad y \in Y,$$

where the function $\phi(y)$ satisfy the equation

$$\phi(u + v) + \phi(u - v) = 2\phi(u) + 2\phi(v), \quad u, v \in Y.$$

Denote by $\Gamma(X)$ the set of all Gaussian distributions on X.

3 Main results

The main result of the article is Theorem 3.2. The prof of Theorem 3.2 uses the following theorem, which is interesting in itself.

Theorem 3.1 *Let X be a discrete Abelian group. Let $\xi_i, i = 1, 2, \ldots, n$, be independent random variables with values in X and distributions μ_i. Consider the linear forms $L_j = \sum_{i=1}^n \alpha_{ij}\xi_i$, where $\alpha_{ij} \in Aut(X), i, j = 1, 2, \ldots, n$. Then the independence of the linear forms L_j implies that $\mu_i = E_{x_i} * m_{G_i}$, where G_i are finite subgroups of X, $x_i \in X$, $i = 1, 2, \ldots, n$.*

Note that Theorem 3.1 was proved earlier in ([3, 13.17]) in the case $n = 2$.

Theorem 3.2 *Let X be a Abelian group of the form*

$$X = \mathbb{R}^m \times K \times D, \tag{2}$$

where, $m \geq 0$ K — compact totally disconnected group, which is topologically isomorphic to a group of the form

$$\mathbf{P}_{p \in \mathfrak{P}}(\triangle_p^{n_p} \times G_p), \tag{3}$$

where n_p are some non negative integers, G_p are finite p-primary groups, D is a discrete countable group. Let $\xi_i, i = 1, 2, \ldots, n$, be independent random variables with values in X and distributions μ_i. Then the independence of the linear forms $L_j = \sum_{i=1}^n \alpha_{ij}\xi_i$, where $\alpha_{ij} \in Aut(X), i, j = 1, 2, \ldots, n$, implies that $\mu_i \in \Gamma(X) * I(X), i = 1, 2, \ldots, n$.

From the conditions of theorem 3.2 it follows that condition (3) is sufficient for validity of the Skitovich-Darmois theorem on compact totally disconnected Abelian groups in the case of n linear forms of n random variables. It turns out, that this condition is necessary.

Theorem 3.3 *Let* X *be a compact totally disconnected Abelian group,* $\xi_i, i = 1, 2, \dots, n$,*be independent random variables with values in* X *and distributions* μ_i, $\alpha_{ij} \in Aut(X)$. *From the independence of the linear forms* $L_j = \sum_{i=1}^{n} \alpha_{ij}\xi_i, j = 1, 2, \dots, n$, *follow that all* $\mu_i \in \Gamma(X) * I(X)$, *if and only if* X *is topologically isomorphic to the group of the form* (3).

Note that the theorem above was proved earlier in ([3, 13.25]) in the case $n = 2$.

Remark. There are locally compact Abelian groups which are non isomorphic to the group of the form (2), and on which The Skitovich-Darmois theorem is valid in the case of n linear forms of n random variables. The needed example was constructed for the case $n = 2$ in [4, 3]. It easily can be checked, that this example can be generalized in the case of the arbitrary n.

References

[1] G. Darmois, Analyse generale des liasions stochastiques. Etude particuliere de l‘analyse factorielle lineaire. *Rev. Inst. Internat. Statistique.* **21** (1953), 2-8.

[2] G.M. Fel'dman, *Arithmetic of Probability Distributions and Characterization Problems on Abelian Groups* AMS translation of mathematical monographs **116**, (American Math. Society, Providence, RI, 1993).

[3] G. Feldman, *Functional equations and Characterizations problems on locally compact Abelian Groups.* (EMS, 2008).

[4] G. M. Feldman and P. Graczyk, The Skitovich-Darmois theorem for locally compact Abelian groups, *J. of the Austral. Math. Soc.* **88** (2010), 339-352.

[5] G. M. Feldman, On the Skitovich-Darmois theorem for finite Abelian groups. *Theory Probabl. Appl.* **37** (1992), 621-631.

[6] G. M. Feldman, On the Skitovich-Darmois theorem on compact groups. *Theory Probab. Appl.* **41** (1996), 768-773.

[7] G. M. Feldman, The Skitovich-Darmois theorem for discrete periodic Abelian groups. *Theory Probab. Appl.* **42** (1997), 611-617.

[8] G. M. Feldman and P. Graczyk, On the Skitovich-Darmois theorem on compact Abelian groups. *J. of Theoretical Probability,* **13** (2000), 859-869.

[9] G. M. Feldman and P. Graczyk, On the Skitovich-Darmois theorem for discrete Abelian groups. *Theory Probab. Appl.* **49** (2005), 527-531.

[10] P. Graczyk and G. M. Feldman, Independent linear statistics on finite Abelian groups. *Ukrainian Math. J.* **53** (2001), 499-506.

[11] A. M. Kagan, Yu. V. Linnik and C. R. Rao, *Characterization Problems in Mathematical Statistics.* Wiley Series in Probability and Mathematical Statistics. (John Wiley & Sons, New York-London-Sydney, 1973).

[12] I. Mazur, The Skitovich-Darmois theorem for finite Abelian groups, arXiv:1106.4134.

[13] K. R. Parthasarathy , R. Ranga Rao, and S. R. S. Varadhan, Probability distributions on locally compact Abelian groups, *Illinois. J. Math.* **7**, 337-369 (1963).

[14] V. P. Skitovich, On a propherty of the normal distribution. *Dokl. Akad. Nauk SSSR (N.S.)* **89** (1953), 217-219.

Mathematical Division
B. Verkin Institute for Low
Temperature Physics and Engineering
of the National Academy
of Sciences of Ukraine
47, Lenin Ave, Kharkov
61103, Ukraine